

References for the Edwards FloTrac system brochure

1. Marik & Cavallazzi. Does central venous pressure predict fluid responsiveness? An updated meta-analysis and a plea for some common sense. *Crit Care Med* 2013
2. Le Manach et al. Can changes in arterial pressure be used to detect changes in cardiac output during volume expansion in the perioperative period? *Anesthesiology* 2013
3. Bennett D. Arterial Pressure: A Personal View. Functional Hemodynamic Monitoring. Berlin: Springer-Verlag, 2005. ISBN: 3-540-22349-5
4. Cecconi M, Fasano N, Langiano N, et al. Goal directed haemodynamic therapy during elective total hip arthroplasty under regional anaesthesia. *Crit Care*. 2011;15:R132
5. Sinclair S, James S, Singer M. Intraoperative intravascular volume optimization and length of hospital stay after repair of proximal femoral fracture: randomised controlled trial. *AMJ*. 1997;315:909–912.
6. Gan T, Soppitt A, Maroof M, et al. Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery. *Anesthesiology*. 2002;97(4):820–826.
7. Venn R, Richardson P, Poloniecki J, Grounds M, Newman P. Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures. *Br J Anaesth*. 2002;88(1):65–71.
8. Conway D, Mayall R, Abdul-Latif M, Gilligan S, Tackaberry C. Randomised controlled trial investigating the influence of intravenous fluid titration using oesophageal Doppler monitoring during bowel surgery. *Anaesthesia*. 2002;57(9):845–849.
9. McKendry M, McGloin H, Saberi D, Caudwell L, Brady A, Singer M. Randomised controlled trial assessing the impact of a nurse delivered, flow monitored protocol for optimisation of circulatory status after cardiac surgery. *BMJ*. 2004;329:358.
10. Wakeling H, McFall M, Jenkins C, Woods W, Barclay G, Fleming S. Intraoperative oesophageal Doppler-guided fluid management shortens postoperative hospital stay after major bowel surgery. *Br J Anaesth*. 2005;95(5):634–642.
11. Noblett S, Snowden C, Shenton B, Horgan A. Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection. *BJS*. 2006;93(9):1069–1076.
12. Chytral, Pradl R, Bosman R, Pelnar P, Kasal E, Zidkova A. Esophageal Doppler-guided fluid management decreases blood lactate levels in multiple-trauma patients: a randomized controlled trial. *Crit Care*. 2007;11:R24.
13. Benes J, Chytral, Altmann P, et al. Intraoperative fluid optimization using stroke volume variation in high risk surgical patients: results of prospective randomized study. *Crit Care*. 2010;14:1-15.
14. Donati A, Loggi S, Preiser JC, Orsetti G, Munch C, Gabbanelli V, Pelaia P, Pietropaoli P. Goal-directed intraoperative therapy reduces morbidity and length of hospital stay in high-risk surgical patients. *Chest*. 2007;132:1817–1824.
15. Grocott et al. Perioperative increase in global blood flow to explicit defined goals and outcomes after surgery: a Cochrane systematic review. *Br J Anaesth* 2013
16. Giglio MT, Marucci M, Testini M, Brienza N. Goal-directed haemodynamic therapy and gastrointestinal complications in major surgery: a meta-analysis of randomized controlled trials. *Br J Anaesth* 2009; 103: 637–66
17. Dalfino L, Giglio MT, Puntillo F, Marucci M, Brienza N. Haemodynamic goal-directed therapy and postoperative infections: earlier is better. A systematic review and meta-analysis. *Crit Care* 2011; 15: R154
18. Corcoran T et al. Perioperative Fluid Management Strategies in Major Surgery: A Stratified Meta-Analysis. *Anesthesia – Analgesia* 2012
19. Hamilton MA, Cecconi M, Rhodes A. A systematic review and meta-analysis on the use of preemptive hemodynamic intervention to improve postoperative outcomes in moderate and high risk surgical patients. *Anesthesia – Analgesia* 2011; 112: 1392–402.
20. Bellamy, MC. Wet, dry or something else? *B J Anaestha*. 2006; 97(6): 755–757

30+ randomized controlled trials confirm benefit

Randomized controlled trials showing a benefit in perioperative goal-directed therapy

21. Shoemaker WC, Appel PL, Kram HB, Waxman K, Lee TS. Prospective trial of supranormal values of survivors as therapeutic goals in high-risk patients. *Chest*. 1988;94(6):1176–1186.
22. Berlauck JF, Abrams, JH, Gilmour, IJ, O'Connor, SR, Knighton, DR, and Cerra, FB. Preoperative optimization of cardiovascular hemodynamics improves outcome in peripheral vascular surgery. A prospective, randomized clinical trial. *Ann Surg*. Sep 1991; 214(3): 289–299.
23. Fleming A, Bishop M, Shoemaker W, Appel P, Sufficool W, Kuvhenguwaha A, Kennedy F, Wo CJ. Prospective trial of supranormal values as goals of resuscitation in severe trauma. *Arch Surg*. 1992 Oct;127(10):1175-9; discussion 1179-81.
24. Boyd O, Grounds RM, Bennett ED. A randomized clinical trial of the effect of deliberate perioperative increase of oxygen delivery on mortality in high-risk surgical patients. *JAMA*. 1993 Dec 8;270(22):2699-707.
25. Mythen MG, Webb AR. Perioperative plasma volume expansion reduces the incidence of gut mucosal hypoperfusion during cardiac surgery. *Arch Surg*. 1995 Apr;130(4):423-9.
26. Sinclair S, James S, Singer M. Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: randomised controlled trial. *BMJ*. 1997 Oct 11;315(7113):909-12.
27. Ueno S, Tanabe G, Yamada H, Kusano C, Yoshidome S, Nuruki K, Yamamoto S, Aikou T. Response of patients with cirrhosis who have undergone partial hepatectomy to treatment aimed at achieving supranormal oxygen delivery and consumption. *Surgery*. 1998 Mar;123(3):278-86.
28. Wilson J, Woods I, Fawcett J, Whall R, Dibb W, Morris C, McManus E. Reducing the risk of major elective surgery: randomised controlled trial of preoperative optimisation of oxygen delivery. *BMJ*. 1999 Apr 24;318(7191):1099-103.
29. Pölonen P, Ruokonen E, Hippeläinen M, Pöyhönen M, Takala J. A prospective, randomized study of goal-oriented hemodynamic therapy in cardiac surgical patients. *Anesth Analg*. 2000 May;90(5):1052-9.
30. Lobo SM, Salgado PF, Castillo VG, Borim AA, Polachini CA, Palchetti JC, Brienzi SL, de Oliveira GG. Effects of maximizing oxygen delivery on morbidity and mortality in high-risk surgical patients. *Crit Care Med*. 2000 Oct;28(10):3396-404.
31. Venn R, Steele A, Richardson P, Poloniecki J, Grounds M, Newman P. Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures. *Br J Anaesth*. 2002 Jan;88(1):65-71.
32. Gan TJ, Soppitt A, Maroof M, El-Moalem H, Robertson KM, Moretti E, Dwane P, and Glass PSA. Goal-directed Intraoperative Fluid Administration Reduces Length of Hospital Stay after Major Surgery. *Anesthesiology*. 2002; 97:820–6.
33. Conway DH, Mayall R, Abdul-Latif MS, Gilligan S, and Tackaberry C. Randomised controlled trial investigating the influence of intravenous fluid titration using oesophageal Doppler monitoring during bowel surgery. *Anaesthesia*. 2002 Sep;57(9):845-9.
34. McKendry M, McGloin H, Saberi D, Caudwell L, Brady AR, and Singer M. Randomised controlled trial assessing the impact of a nurse delivered, flow monitored protocol for optimisation of circulatory status after cardiac surgery. *BMJ*. 2004 Jul 31;329(7460):258. Epub 2004 Jul 8.
35. Wakeling HG, McFall MR, Jenkins CS, Woods WGA, Miles WFA, Barclay GR, and Fleming SC. Intraoperative oesophageal Doppler guided fluid management shortens postoperative hospital stay after major bowel surgery. *BJA*. Volume 95, Issue 5Pp. 634-642.
36. Pearse R, Dawson D, Fawcett J, Rhodes A, Grounds RM, and Bennett ED. Early goal-directed therapy after major surgery reduces complications and duration of hospital stay. A randomised, controlled trial. *Crit Care*. November 2005, 9:R687.
37. Noblett SE, Snowden CP, Shenton BK, and Horgan AF. Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection. *Br J Surg*. 2006 Sep;93(9):1069-76.
38. Chytra I, Pradl R, Bosman R, Pelnar P, Kasal E, and Zidková A. Esophageal Doppler-guided fluid management decreases blood lactate levels in multiple-trauma patients: a randomized controlled trial. *Crit Care*. 2007;11(1):R24.

References continued on back ►



Edwards

39. Lopes MR, Oliveira MA, Pereira VO, Lemos IP, Auler JO Jr, and Michard F. Goal-directed fluid management based on pulse pressure variation monitoring during high-risk surgery: a pilot randomized controlled trial. *Crit Care*. 2007;11(5):R100.
40. Donati A, Loggi S, Preiser JC, Orsetti G, Münch C, Gabbanelli V, Pelaia P, and Pietropaoli P. Goal-directed intraoperative therapy reduces morbidity and length of hospital stay in high-risk surgical patients. *Chest*. 2007 Dec;132(6):1817-24. Epub 2007 Oct 9.
41. Mayer J, Boldt J, Mengistu A, Röhm KD, and Suttner S. Goal-directed intraoperative therapy based on autocalibrated arterial pressure waveform analysis reduces hospital stay in high-risk surgical patients: a randomized, controlled trial. *Crit Care*. 2010; 14:R18.
42. Benes J, Chytra I, Altmann P, Hluchy M, Kasal E, Svitak R, Pradl R, and Stepan M. Intraoperative fluid optimization using stroke volume variation in high risk surgical patients: results of prospective randomized study. *Crit Care*. 2010; 14(3): R118.
43. Jhanji S, Vivian-Smith A, Lucena-Amaro S, Watson D, Hinds CJ, and Pearce RM. Haemodynamic optimisation improves tissue microvascular flow and oxygenation after major surgery: a randomised controlled trial. *Crit Care*. 2010;14(4):R151.
44. Cecconi M, Fasano N, Langiano N, Divella M, Costa MG, Rhodes A, and Rocca GD. Goal-directed haemodynamic therapy during elective total hip arthroplasty under regional anaesthesia. *Crit Care*. 2011; 15:R132.
45. Pillai P, McEleavy I, Gaughan M, Snowden C, Nesbitt I, Durkan G, Johnson M, Cosgrove J, and Thorpe A. A double-blind randomized controlled clinical trial to assess the effect of Doppler optimized intraoperative fluid management on outcome following radical cystectomy. *J Urol*. 2011 Dec;186(6):2201-6.
46. Bisgaard J, Gilsaa T, Ronholm E, and Toft P. Haemodynamic optimisation in lower limb arterial surgery: room for improvement? *Acta Anaesthesiol Scand*. 2013 Feb;57(2):189-98.
47. Ramsingh DS, Sanghvi C, Gamboa J, Cannesson M, Applegate RL 2nd. Outcome impact of goal directed fluid therapy during high risk abdominal surgery in low to moderate risk patients: a randomized controlled trial. *J Clin Monit Comput*. 2013 Jun;27(3):249-57.
48. Zakhaleva J, Tam J, Denoya PI, Bishawi M, and Bergamaschi R. The impact of intravenous fluid administration on complication rates in bowel surgery within an enhanced recovery protocol: a randomized controlled trial. *Colorectal Dis*. 2013 Jul;15(7):892-9.
49. Scheeren TWL, Wiesenack C, Gerlach H, and Marx G. Gernot Marx Goal-directed intraoperative fluid therapy guided by stroke volume and its variation in high-risk surgical patients: a prospective randomized multicentre study. *International Journal of Clinical Monitoring and Computing* (Impact Factor: 1.45). 04/2013.
50. Zhang J, Qiao H, He Z, Wang Y, Che X, and Liang W. Intraoperative fluid management in open gastrointestinal surgery: goal-directed versus restrictive. *Clinics* (Sao Paulo). Oct 2012; 67(10): 1149-1155.
51. Goepfert MS, Richter HP, Zu Eulenburg C, Gruetzmacher J, Rafflenbeul E, Roeher K, von Sandersleben A, Diedrichs S, Reichenspurner H, Goetz AE, and Reuter DA. Individually optimized hemodynamic therapy reduces complications and length of stay in the intensive care unit: a prospective, randomized controlled trial. *Anesthesiology*. 2013 Oct;119(4):824-36.
52. Salzwedel C, Puig J, Carstens A, Bein B, Molnar Z, Kiss K, Hussain A, Belda J, Kirov MY, Sakka SG, and Reuter DA. Perioperative goal-directed hemodynamic therapy based on radial arterial pulse pressure variation and continuous cardiac index trending reduces postoperative complications after major abdominal surgery: a multi-center, prospective, randomized study. *Crit Care*. 2013 Sep 8;17(5):R191.
53. Zheng H, Guo H, Ye JR, Chen L, Ma HP. Goal-directed fluid therapy in gastrointestinal surgery in older coronary heart disease patients: randomized trial. *World J Surg*. 2013 Dec;37(12):2820-9.
54. Zeng K, Li Y, Liang M, Gao Y, Cai H, and Lin C. The influence of goal-directed fluid therapy on the prognosis of elderly patients with hypertension and gastric cancer surgery. *Drug Des Devel Ther*. 2014; 8: 2113-2119.

14 meta-analyses confirm benefit

Hemodynamic optimization through perioperative goal-directed therapy

55. Arulkumaran N, Corredor C, Hamilton MA, et al. Cardiac complications associated with goal-directed therapy in high-risk surgical patients: a meta-analysis. *Br J Anaesth*. 2014 Apr;112(4):648-59.
56. Grocott MP, Dushianthan A, Hamilton MA, et al. Perioperative increase in global blood flow to explicit defined goals and outcomes after surgery: a Cochrane systematic review. *Br J Anaesth*. 2013;111(4):535-548.
57. Aya HD, Cecconi M, Hamilton M, et al. Goal-directed therapy in cardiac surgery: a systematic review and meta-analysis. *Br J Anaesth*, 2013 Apr;110(4):510-7.
58. Cecconi M, Corredor C, Arulkumaran N, et al. Clinical review: Goal-directed therapy - what is the evidence in surgical patients? The effect on different risk groups. *Crit Care Med* 2013; 17:209.
59. Corcoran T, Rhodes JE, Clarke S, et al. Perioperative Fluid Management Strategies in Major Surgery: A Stratified Meta-Analysis. *Anesthesia - Analgesia* 2012; 114(3): 640-651.
60. Dalfino L, Giglio MT, Puntillo F, Marucci M, Brienza N. Haemodynamic goal-directed therapy and postoperative infections: earlier is better. A systematic review and meta-analysis. *Crit Care Med* 2011; 15(3): R154.
61. Hamilton MA, Cecconi M, Rhodes A. A systematic review and meta-analysis on the use of preemptive hemodynamic intervention to improve postoperative outcomes in moderate and high risk surgical patients. *Anesthesia - Analgesia* 2011; 112: 1392-402.
62. Gurgel ST, do Nascimento Jr. P. Maintaining Tissue Perfusion in High-Risk Surgical Patients: A Systematic Review of Randomized Clinical Trials. 2011 *International Anesthesia Research Society*. DOI: 10.1213/ANE.0b013e3182055384.
63. Srinivasa S, Taylor MH, Sammour T, et al. Esophageal Doppler-guided fluid administration in colorectal surgery: critical appraisal of published clinical trials. *Acta Anaesthesiologica Scandinavica* 2011;55(1):4-13.
64. Brienza N, Giglio MT, Marucci M, et al. Does perioperative hemodynamic optimization protect renal function in surgical patients? A meta-analytic study. *Crit Care Med* 2009; 37: 2079-90.
65. Giglio MT, Marucci M, Testini M, et al. Goal-directed haemodynamic therapy and gastrointestinal complications in major surgery: a meta-analysis of randomized controlled trials. *Br J Anaesth*; 2009;103(5):637-646.
66. Phan T, Ismail H, Heriot AG, et al. Improving Perioperative Outcomes: Fluid Optimization with the Esophageal Doppler Monitor, a Metaanalysis and Review. *Journal of the American College of Surgeons*, 2008 Dec;207(6):935-41.
67. Bundgaard-Nielsen M, Holte K, Secher NH, et al. Monitoring of perioperative fluid administration by individualized goal-directed therapy. *Acta Anaesthesiol Scand* 2007 Mar;51(3):331-40.
68. Poeze M, Willem M Greve J, Ramsay G. Meta-analysis of hemodynamic optimization: relationship to methodological quality. *Crit Care* 2005, 9:R771-R779.

For professional use. CAUTION: Federal (United States) law restricts this device to sale by or on the order of a physician. See instructions for use for full prescribing information, including indications, contraindications, warnings, precautions and adverse events.

Edwards Lifesciences devices placed on the European market, meet the essential requirements referred to in Article 3 of the Medical Device Directive 92/42/EEC, and bear the CE marking of conformity.

Edwards, Edwards Lifesciences, the stylized E logo, Clarity In Every Moment, ClearSight, EV1000, FloTrac, Swan-Ganz, and VAMP are trademarks of Edwards Lifesciences Corporation. All other trademarks are the property of their respective owners.

© 2015 Edwards Lifesciences Corporation. All rights reserved. AR12672

Edwards Lifesciences | edwards.com

One Edwards Way | Irvine, California 92614 USA

Switzerland | Japan | China | Brazil | Australia | India



Edwards